

This is a reproduction of a library book that was digitized by Google as part of an ongoing effort to preserve the information in books and make it universally accessible.

GoogleTM books

<https://books.google.com>



Y3.A+7: 22/USNRDL-REPORT

USNRDL Report AD-321(B)
Health and Biology

UNCLASSIFIED

AD-321(B)

**U.S. NAVAL
RADIOLOGICAL DEFENSE
LABORATORY**

SAN FRANCISCO 24, CALIF.



NS 082-001
NM 006-015

THE RELATIONSHIP OF RADIATION DOSE TO LETHALITY
AMONG EXERCISED ANIMALS EXPOSED TO ROENTGEN RAYS

D. J. Kimeldorf and D. C. Jones



UNCLASSIFIED

UNIVERSITY OF MICHIGAN



3 9015 09506 9111

Digitized by Google

This document consists of 18 pages.
This is copy 204 of 300, Series A.

U N C L A S S I F I E D

THE RELATIONSHIP OF RADIATION DOSE TO LETHALITY
AMONG EXERCISED ANIMALS EXPOSED TO ROENTGEN RAYS

D. J. Kimeldorf and D. C. Jones

Interim Report

Technical Objective AW-6

Biological Sciences Branch
C. Entenman, Head

Biological and Medical Sciences Division
M. C. Fishler, Head

U. S. NAVAL RADIOLOGICAL DEFENSE LABORATORY
San Francisco 24, California

Scientific Director
Paul C. Tompkins

11 July 1951

U N C L A S S I F I E D

U N C L A S S I F I E D

DISTRIBUTION

Copies

NAVY

1-4 Chief, Bureau of Ships (Code 348)
 5 Chief, Bureau of Medicine and Surgery (Code 74)
 6 Chief, Bureau of Aeronautics (Code AE54)
 7 Chief, Bureau of Yards and Docks (P-312)
 8-10 Chief, Bureau of Supplies and Accounts (Code OW)
 11 Chief of Naval Research
 12 Office of Naval Research (Galler)
 13 Chief of Naval Operations (Op-36)
 14 Commander, New York Naval Shipyard, Material Laboratory
 15 Director, Naval Research Laboratory
 16 Commanding Officer, Naval Unit, Army Chemical Center
 17 CO, U. S. Naval Civil Engineering, Res. & Eval. Laboratory
 18 U. S. Naval Civil Engineer Corps, Officers School
 19 Naval Medical Research Institute
 20 Chief, Naval Air Experimental Station
 21 Aviation Medical Acceleration Laboratory
 22 Commandant, U. S. Marine Corps
 23 Research and Development Board, Committee on Atomic Energy

ARMY

24 Chief of Engineers, Department of the Army
 25-26 Chief, Research and Engineering Division
 27-28 Office of Chief Signal Officer
 29-30 Office of Chief Signal Officer (Hunt)
 31 President, Army Field Forces Board No. 1, Fort Bragg
 32 CO, Army Medical Research Laboratory
 33 CG, Philadelphia Quartermaster Department
 34 Quartermaster General, Department of the Army
 35 CO, Engineer Research and Development Laboratory
 36 Director, Army Medical Library
 37 Medical Research and Development Board
 38 Army Medical Service Graduate School (Stone)
 39 Radiological Division, CmLC Chemical and Radiological Laboratories
 40-41 CO, CmLC Chemical and Radiological Laboratories
 42 Director of Operations Research (Johnson)
 43 Operations Research Office
 44 Atomic Energy Branch (Betts)

AFSWP

45 Chief, Armed Forces Special Weapons Project
 46 AFSWP, Weapons Defense Division
 47 AFSWP, Sandia Base

U N C L A S S I F I E D

U N C L A S S I F I E D

DISTRIBUTION (Continued)

CopiesAIR FORCE

48-49 Director of Research and Development DCS/D, Hq., USAF (AFDRD-RE)
50 Director of Research and Development (Bruner)
51 CG, Wright Air Development Center (WCRTH-11)
52 CG, Headquarters, Air Materiel Command (MCMTE-11)
53-57 Wright-Patterson Air Force Base (CADO-EL)
58 Wright-Patterson Air Force Base (WRCDO)
59 Air Research and Development Command (DCS/Research)
60 Kirtland Air Force Base (Hooks)
61 Commanding General, Offutt Air Force Base
62 Office of Air Surgeon (Gagge)

ATOMIC ENERGY COMMISSION

63 AEC, Military Applications Division
64 Dr. R. Antonelli
65-72 Argonne National Laboratory
73 Armed Forces Institute of Pathology
74-76 Atomic Energy Commission, Washington
77-80 Atomic Energy Project, Chalk River
81 Battelle Memorial Institute
82 Brush Beryllium Company
83-86 Brookhaven National Laboratory
87-90 Carbide and Carbon Chemicals Division (K-25 Plant)
91-94 Carbide and Carbon Chemicals Division (Y-12 Area)
95 Cleveland Area Office
96 Columbia University (Dunning)
97 Columbia University (Failla)
98-102 duPont Company
103-108 General Electric, Richland
109 Dr. B. Harris
110 Harshaw Chemical Corporation
111-116 Idaho Operations Office
117-118 Iowa State College
119 Isotopes Division (McCormick)
120 Kellex Corporation
121-122 Knolls Atomic Power Laboratory
123-124 Library of Congress, Acquisition Department (Comm)
125-127 Los Alamos Scientific Laboratory
128 Mallinckrodt Chemical Works
129 Massachusetts Institute of Technology (Kaufmann)
130-132 Mound Laboratory
133 National Advisory Committee for Aeronautics
134 National Bureau of Standards (Huntoon)

U N C L A S S I F I E D

U N C L A S S I F I E D
DISTRIBUTION (Continued)

Copies

135	National Bureau of Standards (Library)
136	National Institute of Health
137	National Research Council, Ottawa
138-139	ANP Project
140	Nevis Cyclotron Laboratories
141	New Brunswick Laboratory
142-144	New York Operations Office
145	North American Aviation, Inc.
146	Nucleonics
147-148	Oak Ridge Institute of Nuclear Studies
149-156	Oak Ridge National Laboratory (X-10 Site)
157	Patent Branch, Washington
158	RAND Corporation
159	Sandia Corporation
160	Santa Fe Operations Office
161	Savannah River Operations Office
162	Sylvania Electric Products, Inc.
163-237	Technical Information Service, Oak Ridge
238-247	United Kingdom Scientific Mission (Greenhill)
248	U. S. Geological Survey (Nolan)
249-250	U. S. Public Health Service
251	University of California at Los Angeles
252-257	University of California Radiation Laboratory
258	University of Chicago Toxicity Laboratory
259-260	University of Rochester
261	University of Washington
262	UT-AEC Agricultural Research Program (Hobbs)
263-266	Western Reserve University
267-268	Westinghouse Electric Corporation
 <u>USNRDL</u>	
269-300	USNRDL, Technical Information Division

Date issued: 18 July 1951

U N C L A S S I F I E D

U N C L A S S I F I E D

ABSTRACT

The mortality of exercised irradiated rats was compared with that of non-exercised irradiated rats following x-ray doses ranging from 200 to 1000 r. Swimming to exhaustion under standard conditions of testing constituted the exercise.

Where exposure to x-rays is followed by daily exhaustive exercise the mortality is significantly greater than with irradiation alone. The calculated median lethal dose of x-rays for non-exercised animals is 28 percent higher than that calculated for exercised animals. Several deaths occurred among exercised animals at a radiation dose which was not lethal for non-exercised animals. The mortality is not affected by exercise undertaken prior to irradiation. Experiments demonstrated the fact that the specific environment associated with the method of exercise is not responsible for the mortality increase observed with exercise.

U N C L A S S I F I E D

U N C L A S S I F I E DTHE RELATIONSHIP OF RADIATION DOSE TO LETHALITY
AMONG EXERCISED ANIMALS EXPOSED TO ROENTGEN RAYS*

D. J. Kimeldorf and D. C. Jones

INTRODUCTION

The irradiated mammal appears to be unusually sensitive to a variety of experimental conditions which are tolerated by the non-irradiated animal. Prolonged exposure to cold,¹ daily exhaustive exercise,² exposure to normally non-pathogenic organisms,³ and controlled trauma⁴ are examples of diverse treatment to which irradiated animals are considerably more sensitive than non-irradiated control animals. In addition, these conditions are capable of altering the mortality resulting from exposure to radiation: when irradiated animals are subjected to these conditions the result is a greater number of deaths than is observed following irradiation alone.

The purpose of the present study is to assess the extent to which exhaustive exercise influences the mortality of rats irradiated with various doses of x-rays. Mortality data were obtained on exercised rats exposed to a dose range of x-rays found to be sub-lethal to absolutely lethal for simultaneously irradiated non-exercised control rats. Several additional correlative experiments were carried out in order to analyze the relationships between the conditions of exercise and the altered mortality response observed for exercised irradiated animals.

MATERIALS AND METHODS

Male rats of the Sprague-Dawley strain, bred in the laboratory colony and maintained in air-conditioned quarters under optimum conditions, were used in this study. Animals of comparable age and weight were selected to determine effects at the various x-ray dose levels. All animals were weighed daily for two weeks prior to experimentation and individuals not having normal growth increments were discarded. Of the remainder, only those within \pm eight percent of the mean weight were accepted. The animals in each weight class were from litters born the same week. Food and water were supplied ad lib. except that food was withdrawn two hours prior to each swimming trial. In all studies the animals were caged individually.

* This work was made possible, in part, through funds provided by the Bureau of Medicine and Surgery, U. S. Navy Department.

U N C L A S S I F I E D

U N C L A S S I F I E D

The Exercise Performance Test. The exercised animals performed a standardized exhaustive exercise test developed in this Laboratory for rats. In brief, they were swum individually in 24-gal. cylindrical metal tanks filled with fresh water to a constant level. These tanks were thoroughly cleaned daily and were refilled prior to each day's trials. The water temperature was adjusted to a given temperature $\pm 1.5^{\circ}\text{C}$ within the range of 15° to 21°C . A lead weight was clipped to the chest fur during each swimming trial. Test rats swam until they sank and could no longer rise above a line approximately 18 in. below the surface of the water. The animal was then retrieved and the duration of the swimming trial was recorded. Exhaustive exercise trials were held daily five times per week for two weeks prior to irradiation and for four weeks following irradiation. The pre-irradiation exercise trials were designed to accommodate animals to the test conditions of exercise. After preliminary conditioning for two weeks most animals could swim 15-30 minutes before becoming exhausted under the test conditions.

Irradiation Procedures. The radiation factors for most of the experiments were 250 KVP x-rays; 15 ma; 0.5 mm Cu + 1.0 mm Al (HVL 1.5 mm Cu); 40 in. TSD; 25 r per min. (air dose) measured with Victoreen thimble chambers. In a few studies the dose rate was altered to 8.5 r per min. with the amperage changed to 8 ma. For exposure to the x-ray beam, animals were placed in lucite chambers spaced radially on a motor-driven turntable. The turntable was revolved slowly in the radiation field. Twenty animals were placed in the beam at each exposure.

In most instances the data reported are the cumulative survival data for two or more experiments at each dose of radiation. Only experiments in which irradiated control and experimental animals were under the beam simultaneously are reported. Data were combined from matched experiments since it was felt that repeated studies would increase the reliability of the experimental data. Post-irradiation exercise was initiated on the day of irradiation, which is considered day one in survival data.

RESULTS

In the study of the dose--mortality relationships, 326 non-exercised and 319 exercised irradiated male rats were used. The mortality data resulting from these studies are summarized in Figure 1 and Table 1.

Over the dose range studied, the mortality for exercised animals was greater per roentgen than for non-exercised animals up to the region of the absolute lethal dose (1000 r). The median lethal dose for exercised irradiated animals was significantly lower than that for non-exercised animals. The calculated median lethal dose for non-exercised irradiated animals was 710 r with a probability of 19/20 that the true MLD for these data is

U N C L A S S I F I E D

U N C L A S S I F I E D

between 667 r and 757 r. For exercised irradiated animals the MLD was 510 r with the same probability that the true median lethal dose is between 479 r and 544 r. At the median lethal dose, irradiation followed by daily exhaustive exercise was approximately 1.4 times more potent than irradiation without exercise in the mortality produced. Exhaustive exercise in the absence of radiation was ineffective as a cause of death among the more than 500 non-irradiated exercised animals. Deaths occurred with 400 r when irradiated animals were exercised, while all non-exercised irradiated animals survived this dose. Apparently, daily exhaustive exercise is a stress sufficient to cause deaths when concurrent with a dose of radiation ordinarily not lethal for irradiated animals.

The post-irradiation survival time of exercised animals was altered

U N C L A S S I F I E D

The observed survival for the period of testing is tabulated for purposes of comparison in Table 3. Probability of a difference as great as that observed between the irradiated controls and experimental groups occurring by chance is indicated. These probability values were obtained by the corrected chi-square formula method.

Exposure to water at either temperature did not alter the mortality of irradiated animals significantly from that of the non-exposed irradiated control animals. The survival rate for irradiated animals exposed to 15°C water did not differ from that of animals exposed to 21°C water ($p > 0.50$). Neither the nature nor the degree of an exposure stress was detectable by body weight changes or by clinical observations of the water-exposed animals. The survival of irradiated animals exercised in water of either temperature was significantly lower than that of non-exercised irradiated controls ($p < 0.01$), and the mortality rates did not differ significantly between the irradiated animals exercised at the two temperatures ($p > 0.10$). It appears, then, that the irradiation mortality is relatively independent of the water temperature (over the range tested) but is dependent upon whether the animals were or were not exercised following irradiation. In addition, merely exposing the irradiated rats to water after irradiation has no significant effect upon mortality.

An additional study was undertaken to determine if exposure to water could alter the length of the post-irradiation survival period. A highly lethal radiation dose (860 r) was used in order to provide the experiment with large numbers of animals dying during the observation period. Approximately 190 animals were split into groups as designated in Table 4. The duration and conditions of exposure have been previously described. Animals were exposed to water daily five days per week for two weeks prior to irradiation; after irradiation they were exposed daily five times per week until the last experimental animal had died. The effective time for 50 percent mortality was calculated by Litchfield's probit method.⁶ Data for a series of exercised animals irradiated with 860 r are included for purposes of comparison in Table 4.

The median survival times were nearly identical for all water-exposed irradiated groups. It is concluded that exposure to water does not, by itself, significantly alter the post-irradiation survival time after highly lethal amounts of radiation.

Animals used in the principal studies were accommodated to the swimming tests by daily exhaustive exercise five times per week for two weeks prior to irradiation. The influence of this pre-irradiation exercise upon the mortality during the post-irradiation period was studied in 168 animals. These were segregated into groups as tested and summarized in Table 5.

U N C L A S S I F I E D

U N C L A S S I F I E D

It is apparent in this experiment that the survival of irradiated animals is significantly altered by daily exhaustive exercise after irradiation, but not by exercise prior to radiation exposure. Accommodation to the test by exercise prior to irradiation did not markedly influence the mortality of animals exercised after irradiation ($p > 0.20$).

DISCUSSION

As measured by the resulting mortality, the irradiated animal is highly sensitive to the stress of daily exhaustive exercise in the swimming performance test. The extent of this stress is somewhat indicated by the occurrence of deaths (23 percent) among exercised animals following an amount of radiation (400 r) which is non-lethal for normal animals. In order to alter the mortality, it is deduced that exercise must occur after irradiation, since exercise prior to irradiation was shown to not significantly influence mortality. Since the probit curves for mortality of exercised and non-exercised animals are parallel, within the experimental error, it appears that the influence of exercise upon mortality is a function of the mortality level which occurs with non-exercised animals in the dose range studied. This suggests that the mechanism of mortality is similar for exercised and non-exercised animals.

Swimming to exhaustion was selected as the technique of exercise in these studies since it represents an intensive form of muscular effort during high motivation, presumably with survival as the goal. This method has been used in several studies of the physiological state associated with intense muscular activity^{7,8,9} and in studies evaluating the capacity to do work.¹⁰⁻¹³ It is recognized that the swimming procedure necessitates exposure of animals to an unusual environment for a brief period daily. In view of the mortality data obtained from the studies of water exposure, it is concluded that exposure to the conditions of testing is not responsible for the mortality observed with swum-irradiated animals.

Smith and Smith¹⁴ were unable to alter significantly the mortality rate of irradiated mice by subjecting them to forced exercise on a treadmill. Under the conditions of the experiment, the mice were exercised for a specified duration of time, although not necessarily to exhaustion. From the data of Steinhaus and his collaborators^{15,16} it appears that the intensity of exercise occurring during swimming performance is greater than that obtained by treadmill techniques in terms of physiological response. In view of the results obtained by Smith and Smith, it may be that intensity of exercise and degree of exhaustion are the significant factors altering the mortality of irradiated animals by exercise. This assumes that there are no species differences involved. These factors are being varied independently in performance studies now being pursued at this Laboratory.

U N C L A S S I F I E D

U N C L A S S I F I E D

SUMMARY

If animals are exercised by swimming to exhaustion daily after irradiation, there results a greater mortality, with more deaths at lower doses, than with irradiation alone. The calculated median lethal dose of x-rays for non-exercised animals is 28 percent higher than that calculated for exercised animals. Exercise effectively alters the level of mortality when undertaken after irradiation, whereas exercise prior to irradiation has no significant effect upon mortality.

Approved by: *McFishler*
Head, Biological and
Medical Sciences Division

For the Scientific Director

Acknowledgments

The authors are grateful for the advice and criticism of Dr. M. C. Fishler and the technical assistance of T. J. Castanera, G. K. Osborn, and J. L. Lee during several phases of this study.

U N C L A S S I F I E D

U N C L A S S I F I E D

REFERENCES

1. Smith, W. W., B. J. Highman, J. R. Mitchell and H. C. Blount, Jr. Proc. Soc. Exper. Biol. & Med. 71:498, 1949.
2. Kimeldorf, D. J., D. C. Jones and M. C. Fishler. Science. 112:175, 1950.
3. Schechmeister, I. L. and V. P. Bond. U. S. Naval Radiological Defense Lab. Report AD-233(B), Aug. 11, 1950. San Francisco, California.
4. Chapman, W. H., C. R. Sipe, D. C. Eltzholtz, E. P. Cronkite and F. W. Chambers. Radiology. 55:865, 1950.
5. Litchfield, J. T., Jr., and F. Wilcoxon. J. of Pharm. & Exper. Ther. 95:99, 1949.
6. Litchfield, J. T., Jr. J. of Pharm. & Exper. Therap. 97:399, 1949.
7. Steinhaus, A. H. Physiol. Rev. 13:103, 1933.
8. Miller, H. C. and D. C. Darrow. Am. J. Physiol. 132:801, 1941.
9. Postel, S., J. M. Tobias, H. M. Patt and R. W. Gerard. Proc. Soc. Exper. Biol. & Med. 63:432, 1946.
10. Harris, R. and D. Ingle. Psychol. Bull. 35:667, 1938.
11. Ramey, E., M. S. Goldstein and R. Levine. Am. J. Physiol. 162:10, 1950.
12. Scheer, B. T., S. Dorst, J. F. Codie and D. F. Soule. Am. J. Physiol. 149:194, 1947.
13. Gaarenstroom, J., L. Waterman and E. Laquer. Acta Brevia Neerl. Physiol. 7:10, 1936.
14. Smith, F. and W. W. Smith. Federation Proc. 9:117, 1950.
15. Steinhaus, A. H., L. A. Hoyt and H. A. Rice. Am. J. Physiol. 99:521, 1931.
16. Rice, H. A. and A. H. Steinhaus. Am. J. Physiol. 96:592, 1930.

U N C L A S S I F I E D

U N C L A S S I F I E D

TABLE 1

Analysis of mortality data by the probit methods of Litchfield and Wilcoxon.⁵

Experimental Conditions	No. of Animals	LD ₅₀ (r)	Confidence Limits LD ₅₀ (p = 0.95)	Slope Function	Confidence Limits of Slope Function (p = 0.95)
Non-exercised irradiated (200-1000 r)	326	710	667-757	1.23	1.17-1.29
Exercised irradiated (200-1000 r)	319	510	479-544	1.29	1.24-1.33
Experimental condition tested	SR/f _{SR} *		Confidence limits SR (p = 0.95)	PR/f _{PR} **	Confidence limits PR (p = 0.95)
Exercise	1.05/1.07		0.98-1.13	1.39/1.08	1.29-1.50

* If SR (slope ratio) is less than f_{SR}, then the experimental condition is comparable with the control condition within experimental error.

** If PR (potency ratio) exceeds f_{PR}, then there is a significant shift in the degree of lethality attributable to exercise.

U N C L A S S I F I E D

U N C L A S S I F I E D

TABLE 2

Distribution of deaths for animals dying during the 30 days of observation. The cumulative mortality is indicated for seven day intervals.

Dose	Conditions	Number of Animals	First Death (Days PI*)	Percent Dead				Median Time Survival (Days PI*)
				Days 1-7	Days 8-14	Days 15-21	Days 22-28	
200 r	Non-Exercised	20	-		None died			---
	Exercised	19	-		None died			---
300 r	Non-Exercised	20	-		None died			---
	Exercised	16	-		None died			---
400 r	Non-Exercised	60	-	0	0	0	0	---
	Exercised	43	13	0	7	12	23	18
500 r	Non-Exercised	84	8	0	5	5	5	12
	Exercised	77	5	5	17	36	40	14
600 r	Non-Exercised	40	6	8	22	25	28	9
	Exercised	60	8	0	23	52	68	15
700 r	Non-Exercised	38	8	0	21	44	44	13
	Exercised	36	3	14	81	92	92	10
860 r	Non-Exercised	40	5	20	88	88	88	9
	Exercised	43	2	84	98	100	100	4
1000 r	Non-Exercised	24	4	100	100	100	100	4
	Exercised	25	3	100	100	100	100	4

* Days post-irradiation.

U N C L A S S I F I E D

U N C L A S S I F I E D

TABLE 3

Percent survival for irradiated animals (500 r) following daily treatment as noted for two weeks prior to irradiation, and for four weeks post-irradiation.

Experimental Condition	No. of Animals	No. Alive	Percent Alive	p
Irradiated controls	65	60	92	
Exposed to 15°C water	25	22	88	>0.80
Exposed to 21°C water	25	24	94	>0.80
Exercised in 15°C water	42	30	71	<0.01
Exercised in 21°C water	54	29	54	<0.01
Non-irrad. exposed (15°C)	25	25	100	>0.30
Non-irrad. exercised (15°C)	25	25	100	>0.30

U N C L A S S I F I E D

U N C L A S S I F I E D

TABLE 4

The median survival time (ET-50%) for groups of irradiated animals (860 r) exposed to water but not exercised. The ET-50% was calculated by the probit method of Litchfield.⁶

Experimental Conditions	Animals	Mortality	ET-50% (Days)	Confidence Interval (p = 0.95)
Irradiated controls	37	100	9.0	8.6-9.4
Exposed prior to irradiation	39	100	9.5	8.8-10.3
Exposed after irradiation	37	100	9.3	8.9-9.7
Exposed before and after irradiation	39	100	8.6	8.1-9.1
Exercised before and after irradiation	43	100	4.3	3.8-4.8
Exposed but not irradiated	33	0	---	-----

TABLE 5

Percent survival for irradiated animals (500 r) following daily exercise for two weeks prior to irradiation and/or four weeks post-irradiation.

Experimental Conditions	No. of Animals	No. Alive	Percent Alive	p
Irradiated only	40	39	97	
Exercised prior to irradiation only	35	33	94	>0.80
Exercised after irradiation only	31	24	77	<0.05
Exercised before and after irradiation	62	40	64	<0.01

U N C L A S S I F I E D

UNCLASSIFIED




FIG. 1 Percent survival of exercised and non-exercised irradiated rats at 30 days post-irradiation (X-ray).

UNCLASSIFIED

UNCLASSIFIED